



May 30, 2019

Dear 7<sup>th</sup> grade parents,

### Vaccine requirements for 7<sup>th</sup> graders

The state of Ohio requires all students entering the 7<sup>th</sup> grade to have the meningococcal vaccine (also known as MCV4, Menevo or Menactra) in addition to one dose of Tdap (Tetanus, diphtheria and pertussis).

#### **Meningococcal meningitis**

Meningococcal meningitis is a rare but serious disease that develops rapidly and can claim a life in as little as one day. Of those who survive, approximately one in five are left with serious medical problems like amputation, deafness, and brain damage. Teens are at increased risk of meningococcal meningitis. This increased risk may be due to activities like sharing utensils and kissing.

#### **Tetanus, diphtheria, and pertussis (Tdap)**

Tetanus causes painful tightening of the muscles, usually all over your body; diphtheria causes a thick covering in the back of the throat and can also lead to breathing problems, paralysis, heart failure, and even death; pertussis causes coughing spells and can lead to pneumonia, seizures, brain damage, and death, particularly in infants.

Vaccination is the best way to protect someone from getting the disease.

You may obtain these vaccines from your child's health care provider or other community resource. Vaccines are also available at your local health department. See attached flyer for health department open hours.

Students who do not provide documentation of these immunizations to the school this fall are subject to exclusion. If there are medical/religious or philosophical reasons your child cannot receive immunizations, please contact the school clinic for an Immunization Exemption Form.

Thank you,

Jennifer Henderson, MSN, RN, NCSN  
District RN Supervisor for Copley-Fairlawn City Schools  
Akron Children's Hospital, School Health Services

*Attached CDC "2019 Recommended Immunizations for Children from 7 Through 18 Years Old"*

## INFORMATION FOR PARENTS

## 2019 Recommended Immunizations for Children 7–18 Years Old

### Talk to your child's doctor or nurse about the vaccines recommended for their age.

|                          | Flu Influenza  | Tdap Tetanus, diphtheria, pertussis                       | HPV Human papillomavirus  | Meningococcal   | Hepatitis B  | Hepatitis A  | Polio | MMR Measles, mumps, rubella | Chickenpox Varicella |
|--------------------------|--|---|---|---|--|--------------|-------|-----------------------------|----------------------|
|                          |  |   |   | MenACWY   | MenB   | Pneumococcal |       |                             |                      |
| <b>7-8 Years</b>         |  |   |   |   |  |              |       |                             |                      |
| <b>9-10 Years</b>        |  |   |   |   |  |              |       |                             |                      |
| <b>11-12 Years</b>       |  |   |   |   |  |              |       |                             |                      |
| <b>13-15 Years</b>       |  |   |   |   |  |              |       |                             |                      |
| <b>16-18 Years</b>       |  |   |   |   |  |              |       |                             |                      |
| <b>More information:</b> | Everyone 6 months and older should get a flu vaccine every year. | All 11- through 12-year olds should get one shot of Tdap. | All 11- through 12-year olds should get a 2-shot series of HPV vaccine. A 3-shot series is needed for those with weakened immune systems and those who start the series at 15 years or older. | All 11- through 12-year olds should get one shot of meningococcal conjugate (MenACWY). A booster shot is recommended at age 16. | Teens 16-18 years old may be vaccinated with a serogroup B meningococcal (MenB) vaccine. |              |       |                             |                      |


These shaded boxes indicate when the vaccine is recommended for all children unless your doctor tells you that your child cannot safely receive the vaccine.

These shaded boxes indicate the vaccine should be given if a child is catching up on missed vaccines.

This shaded box indicates children not at increased risk may get the vaccine if they wish after speaking to a provider.



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention



American Academy  
of Pediatrics  
DEDICATED TO THE HEALTH OF ALL CHILDREN®  
AMERICAN ACADEMY OF FAMILY PHYSICIANS

AAFP  
AMERICAN ACADEMY OF FAMILY PHYSICIANS

AAFP  
AMERICAN ACADEMY OF FAMILY PHYSICIANS

# Vaccine-Preventable Diseases and the Vaccines that Prevent Them

## Diphtheria (Can be prevented by Tdap vaccination)

Diphtheria is a very contagious bacterial disease that affects the respiratory system, including the lungs. Diphtheria bacteria can be spread from person to person by direct contact with droplets from an infected person's cough or sneeze. When people are infected, the bacteria can produce a toxin (poison) in the body that can cause a thick coating in the back of the nose or throat that makes it hard to breathe or swallow. Effects from this toxin can also lead to swelling of the heart muscle and, in some cases, heart failure. In serious cases, the illness can cause coma, paralysis, or even death.

## Hepatitis A (Can be prevented by HepA vaccination)

Hepatitis A is an infection in the liver caused by hepatitis A virus. The virus is spread primarily person to person through the fecal-oral route. In other words, the virus is taken in by mouth from contact with objects, food or drinks contaminated by the feces (stool) of an infected person. Symptoms can include fever, tiredness, poor appetite, vomiting, stomach pain, and sometimes jaundice (when skin and eyes turn yellow). An infected person may have no symptoms, may have mild illness for a week or two, may have severe illness for several months, or may rarely develop liver failure and die from the infection. In the U.S., about 100 people a year die from hepatitis A.

## Hepatitis B (Can be prevented by HepB vaccination)

Hepatitis B causes a flu-like illness with loss of appetite, nausea, vomiting, rashes, joint pain, and jaundice. Symptoms of acute hepatitis B include fever, fatigue, loss of appetite, nausea, vomiting, pain in joints and stomach, dark urine, grey-colored stools, and jaundice (when skin and eyes turn yellow).

## Human Papillomavirus (Can be prevented by HPV vaccination)

Human papillomavirus is a common virus. HPV is most common in people in their teens and early 20s. About 14 million people, including teens, become infected with HPV each year. HPV infection can cause cervical, vaginal, and vulvar cancers in women and penile cancer in men. HPV can also cause anal cancer, oropharyngeal cancer (back of the throat), and genital warts in both men and women.

## Influenza (Can be prevented by annual flu vaccination)

Influenza is a highly contagious viral infection of the nose, throat, and lungs. The virus spreads easily through droplets when an infected person coughs or sneezes and can cause mild to severe illness. Typical symptoms include a sudden high fever, chills, a dry cough, headache, runny nose, sore throat, and muscle and joint pain. Extreme fatigue can last from several days to weeks. Influenza may lead to hospitalization or even death, even among previously healthy children.

## Measles (Can be prevented by MMR vaccination)

Measles is one of the most contagious viral diseases. Measles virus is spread by direct contact with the airborne respiratory droplets of an infected person. Measles is so contagious that just being in the same room after a person who has measles has already left can result in infection. Symptoms usually include a rash, fever, cough, and red, watery eyes. Fever can persist, rash can last for up to a week, and coughing can last about 10 days. Measles can also cause pneumonia, seizures, brain damage, or death.

## Meningococcal Disease (Can be prevented by meningococcal vaccination)

Meningococcal disease has two common outcomes: meningitis (infection of the lining of the brain and spinal cord) and bloodstream infections. The bacteria that cause meningococcal disease spread through the exchange of nose and throat droplets, such as when coughing, sneezing, or kissing. Symptoms include sudden onset of fever, headache, and stiff neck. With bloodstream infection, symptoms also include a dark purple rash. About one of every 10 people who gets the disease dies from it. Survivors of meningococcal disease may lose their arms or legs, become deaf, have problems with their nervous systems, become developmentally disabled, or suffer seizures or strokes.

## Mumps (Can be prevented by MMR vaccination)

Mumps is an infectious disease caused by the mumps virus, which is spread in the air by a cough or sneeze from an infected person. A child can also get infected with mumps by coming in contact with a contaminated object like a toy. The mumps virus causes swollen salivary glands under the ears or jaw, fever, muscle aches, tiredness, abdominal pain, and loss of appetite. Severe complications for children who get mumps are uncommon, but can include meningitis (inflammation of the lining of the brain and spinal cord), encephalitis (inflammation of the brain), permanent hearing loss, or swelling of the testes, which rarely results in decreased fertility.

## Pertussis (Whooping Cough) (Can be prevented by Tdap vaccination)

Pertussis spreads very easily through coughing and sneezing. It can cause a bad cough that makes someone gasp for air after coughing fits. This cough can last for many weeks, which can make preteens and teens miss school and other activities. Pertussis can be deadly for babies who are too young to receive the vaccine. Often babies get whooping cough from their older brothers or sisters, like preteens or teens, or other people in the family. Babies with pertussis can get pneumonia, have seizures, become brain damaged, or even die. About half of children under 1 year of age who get pertussis must be hospitalized.

## Pneumococcal Disease (Can be prevented by pneumococcal vaccination)

Pneumonia is an infection of the lungs that can be caused by the bacteria called "pneumococcus." These bacteria can cause other types of infections, too, such as ear infections, sinus infections, meningitis (infection of the lining of the brain and spinal cord), and bloodstream infections. Sinus and ear infections are usually mild and are much more common than the more serious forms of pneumococcal disease. However, in some cases, pneumococcal disease can be fatal or result in long-term problems like brain damage and hearing loss. The bacteria that cause pneumococcal disease spread when people cough or sneeze. Many people have the bacteria in their nose or throat at one time or another without being ill—this is known as being a carrier.

## Polio (Can be prevented by IPV vaccination)

Polio is caused by a virus that lives in an infected person's throat and intestines. It spreads through contact with the stool of an infected person and through droplets from a sneeze or cough. Symptoms typically include sore throat, fever, tiredness, nausea, headache, or stomach pain. In about 1% of cases, polio can cause paralysis. Among those who are paralyzed, about 2 to 10 children out of 100 die because the virus affects the muscles that help them breathe.

## Rubella (German Measles) (Can be prevented by MMR vaccination)

Rubella is caused by a virus that is spread through coughing and sneezing. In children, rubella usually causes a mild illness with fever, swollen glands, and a rash that lasts about 3 days. Rubella rarely causes serious illness or complications in children, but can be very serious to a baby in the womb. If a pregnant woman is infected, the result for the baby can be devastating, including miscarriage, serious heart defects, mental retardation, and loss of hearing and eyesight.

## Tetanus (Lockjaw) (Can be prevented by Tdap vaccination)

Tetanus mainly affects the neck and belly. When people are infected, the bacteria produce a toxin (poison) that causes muscles to become tight, which is very painful. This can lead to "flocking" of the jaw so a person cannot open his or her mouth, swallow, or breathe. The bacteria that cause tetanus are found in soil, dust, and manure. The bacteria enter the body through a puncture, cut, or sore on the skin. Complete recovery from tetanus can take months. One to two out of 10 people who get tetanus die from the disease.

## Varicella (Chickenpox) (Can be prevented by varicella vaccination)

Chickenpox is caused by the varicella zoster virus. Chickenpox is very contagious and spreads very easily from infected people. The virus can spread from either a cough or sneeze. It can also spread from the blisters on the skin, either by touching them or by breathing in these viral particles. Typical symptoms of chickenpox include an itchy rash with blisters, tiredness, headache, and fever. Chickenpox is usually mild, but it can lead to severe skin infections, pneumonia, encephalitis (brain swelling), or even death.

If you have any questions about your child's vaccines, talk to your child's doctor or nurse.